



TITLE:

2-Microlocal Boundary Value Problems and Their Applications(Functional-Analytic Study of Generalized Functions)

AUTHOR(S):

Uchida, Motoo

CITATION:

Uchida, Motoo. 2-Microlocal Boundary Value Problems and Their Applications(Functional-Analytic Study of Generalized Functions). 数理解析研究所講究録 1989, 704: 39-42

ISSUE DATE:

1989-10

URL:

<http://hdl.handle.net/2433/101595>

RIGHT:

2-Microlocal Boundary Value Problems and Their Applications

by Motoo Uchida (Univ. of Tokyo)

(内田素夫)

1. Let M be a real analytic manifold, X a complex neighbourhood of M , and Y a complex hypersurface of X . Let Λ be a regular involutive conic submanifold of $T^*_M X$, Λ^C (resp. $\tilde{\Lambda}$) a complexification (resp. a partial complexification) of Λ in T^*X . Let \mathfrak{M} be a coherent \mathcal{E}_X -Module defined in a neighbourhood of Λ^C and assume that $Y \rightarrow X$ is non microcharacteristic along Λ^C for \mathfrak{M} (cf. Def. 2.2 of [1] and Def. 2.10.3 of [5]). Set $\Sigma = \Lambda \cap \pi^{-1}(Y)$, and denote by Λ_+ a domain of Λ with boundary Σ . Then we can define the microlocal boundary values along Λ to Σ for \mathcal{B}^2_{Λ} -solutions to \mathfrak{M} . To be precise, there exists the boundary value map

$$bv : R\mathcal{H}om_{\mathcal{E}_X}(\mathfrak{M}, \Gamma_{\Lambda_+}(\mathcal{B}^2_{\Lambda}))|_{\Sigma} \rightarrow R\mathcal{H}om_{\mathcal{E}_Y}(\mathfrak{M}_Y, \mathcal{B}^2_{\Sigma}),$$

where \mathcal{B}^2_{Λ} denotes the sheaf of 2-hyperfunctions on Λ due to Kashiwara (cf. [4], [10]), and \mathfrak{M}_Y denotes the tangential system on Y of \mathfrak{M} .

We set

$$\mathcal{E}^2_{\Lambda_+}|_{\tilde{\Lambda}} = \mu\text{hom}(\mathcal{E}_{\Lambda_+}, \mathcal{E}^h_{\tilde{\Lambda}}) \otimes \text{or}_{\Lambda|\tilde{\Lambda}}[\text{codim } \Lambda],$$

where $\mathcal{E}^h_{\tilde{\Lambda}}$ denotes the sheaf of microfunctions with holomorphic parameters on $\tilde{\Lambda}$, and $\text{or}_{\Lambda|\tilde{\Lambda}}$ denotes the relative orientation sheaf on Λ .

Refer to Kashiwara-Schapira [3] for the bifunctor μhom . This complex of \mathcal{E}_X -Modules is a 2-microlocal version of $\mathcal{E}_{\Omega|X}$, the "sheaf" of microfunctions for boundary value problems, which is introduced in Schapira [8]. This allows us to define the "boundary 2-analytic wavefront set" $SS^2_{\Lambda_+}$ for sections of $\Gamma_{\Lambda_+}(\mathcal{B}^2_{\Lambda})$, and we can analyse the boundary value map "bv" 2-microlocally in terms of $SS^2_{\Lambda_+}$. In particular, we can show the reflection of 2-microlocal singularity at the boundary. Refer to [13] for the details.

2. Let Ω be an open subset of M with real analytic boundary $N = \{\varphi=0\}$, Y the complexification of N in X . We denote by ρ the natural

projection $Y \times_X T^*X \longrightarrow T^*Y$. Take a point $y^* \in T^*_N Y$ and a point $x^* \in T^*_M X \times N$ with $\rho(x^*) = y^*$. Let P be a microdifferential operator defined in a neighbourhood of x^* with involutive double characteristics. Precisely we assume that the principal symbol $\sigma(P)$ of P is decomposed by homogeneous holomorphic functions p_1, p_2, q :

$$\sigma(P) = q \cdot p_1^{m_1} \cdot p_2^{m_2}$$

in a neighbourhood of x^* and they satisfy the following conditions:

- (1) p_1 and p_2 are real valued on $T^*_M X$,
- (2) $p_1(x^*) = p_2(x^*) = 0$, $q(x^*) \neq 0$,
- (3) $dp_1 \wedge dp_2 \wedge \omega \neq 0$,
- (4) $\{p_1, p_2\} = 0$ on $\Lambda = \{p_1 = p_2 = 0\}$,
- (5) $\{q, p_i\} \neq 0$ ($i=1, 2$).

In this situation we consider the microlocal boundary value problem

$$(D) \quad \begin{cases} Pu = 0 \text{ at } x^*, \\ (\varphi_X \cdot D_X)^i u|_{\varphi \rightarrow +0} = 0 \text{ at } y^* \quad (0 \leq i < \max(m_1, m_2)). \end{cases}$$

Remark that we take here the boundary value $(\varphi_X \cdot D_X)^i u|_{\varphi \rightarrow +0}$ of u in the microlocal sense from a neighbourhood of x^* .

Then we have the following results.

We denote by $SS_\Omega(u)$ the boundary analytic wavefront set of u (cf. [8] for the definition of the boundary analytic wavefront set).

Theorem 1.---Let Γ be a real bicharacteristic leaf of Λ passing through x^* . For any solution u of the microlocal boundary value problem (D), there exists a subset $\{x^*_s\}$ of $\Gamma \cap \Sigma$ such that


$$SS_\Omega(u) \cap \Gamma = \text{the closure in } \Gamma \text{ of the union of } \{b^i_s; s, i=1, 2\} \text{ and}$$

$$\text{some of connected components of } \Gamma_+ \setminus \bigcup \{b^i_s; s, i=1, 2\},$$

where $\Gamma_+ = \Gamma \times_M \Omega$, and b^i_s denotes the half integral curve of H_{p_i} , the Hamilton vector field of p_i , issued from x^*_s into Γ_+ .

Corollary 2.---For any solution u of the microlocal boundary value problem (D),

$$\begin{aligned} & b^1(x^*) \cup b^2(x^*) \not\subset SS(u|_{\Omega}) \\ \Rightarrow & x^* \notin SS_{\Omega}(u). \end{aligned}$$



Theorem 1 is obtained as an application of the theory of Section 1 (cf. [13]).

Remark.---As for the results in the interior domain for the same operator we refer to Tose [10, 11, 12]. We also refer to Lascar [6] for the similar result as Corollary 2 in the C^∞ category. Note that we assume in Corollary 2 that at least one of the integral curves $b^1(x^*)$, $b^2(x^*)$ is not contained in $SS(u|_{\Omega})$ in a neighbourhood of x^* .

References

- [1] Kashiwara M. and P. Schapira, Problèmes de Cauchy pour les systèmes microdifférentiels dans le domaine complexe. *Invent. Math.* **46**, 17-38 (1978).
- [2] ---, Micro-hyperbolic systems. *Acta Math.* **142**, 1-55 (1979).
- [3] ---, Microlocal study of sheaves. *Asterisque* 128, Soc. Math. France (1985).
- [4] Kashiwara M. and Y. Laurent, Théorèmes d'annulation et deuxième microlocalisation. Prépublication d'Orsay (1983).
- [5] Laurent, Y., Théorie de la deuxième microlocalisation dans le domaine complexe. *Progress in Math.* 53, Birkhauser (1985).
- [6] Lascar, R., Paramétrices microlocales de problèmes aux limites pour des équations pseudo-différentielles à caractéristiques de multiplicité variable. *C. R. Acad. Sc. Paris* **287**, 441-443 (1978).
- [7] Schapira, P., Propagation at the boundary and reflection of analytic singularities of solutions of linear partial differential equations I. *Publ. RIMS Kyoto Univ.* **12** Suppl., 441-453 (1977).
- [8] ---, Front d'onde analytique au bord I, II. *C. R. Acad. Sc. Paris* **302**, 383-386 (1986), *Sem. E.D.P. Ecole Polyt. Exp.* 13 (1986).
- [9] Schapira, P. and G. Zampieri, *Coll. Intern. "Hyperbolic Eq."*, Padova, 1985.
- [10] Tose, N., On a class of microdifferential equations with involutive double characteristics. *J. Fac. Sci. Univ. of Tokyo* **33**, 619-634 (1986).
- [11] ---, The 2-microlocal canonical form for a class of microdifferential equations and propagation of singularities. *Publ. RIMS Kyoto Univ.* **23**, 101-116 (1987).
- [12] ---, On a class of 2-microhyperbolic systemes. *J. Math. pures et appl.* **67**, 23-37 (1988).
- [13] Uchida, M., 2-Microlocal boundary value problems and their applications. Master Thesis, Univ. of Tokyo (1987), (in Japanese).